

Self-similar Bianchi models: II. Class B models

Pantelis S. Apostolopoulos[†]

Abstract. In a companion article (referred hereafter as paper I) a detailed study of the simply transitive Spatially Homogeneous (SH) models of class A concerning the existence of a simply transitive similarity group has been given. The present work (paper II) continues and completes the above study by considering the remaining set of class B models. Following the procedure of paper I we find all SH models of class B subjected only to the minimal geometric assumption to admit a proper Homothetic Vector Field (HVF). The physical implications of the obtained geometric results are studied by specialising our considerations to the case of vacuum and γ -law perfect fluid models. As a result we regain all the known exact solutions regarding vacuum and non-tilted perfect fluid models. In the case of tilted fluids we find the *general* self-similar solution for the exceptional type VI_{-1/9} model and we identify it as equilibrium point in the corresponding dynamical state space. It is found that this *new* exact solution belongs to the subclass of models $n_\alpha^\alpha = 0$, is defined for $\gamma \in (\frac{4}{3}, \frac{3}{2})$ and although has a five dimensional stable manifold there exist always two unstable modes in the restricted state space. Furthermore the analysis of the remaining types, guarantees that tilted perfect fluid models of types III, IV, V and VII_h cannot admit a proper HVF strongly suggesting that these models either may not be asymptotically self-similar (type V) or may be extreme tilted at late times. Finally for each Bianchi type, we give the extreme tilted equilibrium points of their state space.

[†]University of Athens, Department of Physics, Nuclear and Particle Physics Section,
Panepistimiopolis, Zografos 15771, Athens, Greece.

E-mail: papost@phys.uoa.gr

PACS: 04.20.Ha, 04.20.Jb, 04.20.Nr, 98.80.-k

1. Introduction

In a previous article [1] (designated in the following as paper I) we presented a general study of the Spatially Homogeneous (SH) models of class A admitting a four dimensional group of homotheties acting simply transitively on the spacetime manifold. We concentrated on tilted γ -law perfect fluids and we have given various results concerning the existence of an admissible self-similar perfect fluid model in all the Bianchi types of class A. As a consequence this analysis has led us to regain the type II solution [2], to show the non-existence of self-similar type VII₀, VIII and IX tilted perfect fluid models [1, 3] and to prove the generality of the known perfect fluid solution for type VI₀ models [4, 5].

Our intention in the present work is to address the problem regarding the existence of a proper Homothetic Vector Field (HVF) for the remaining SH models namely those of class B and to examine the physical and dynamical implications of the geometric results with the hope to provide a way of obtaining a better understanding of their past and future asymptotic dynamics.

Section 2 contains some basic elements of the two major frameworks of studying SH cosmologies: the *orthonormal frame* and *metric* approaches. In particular and adopting the conventions and the methodology of [6], we briefly discuss the structure of the dynamical state space of the SH models, coming from the evolution equations and the algebraic restrictions for the case where the spacetime is filled with a tilted (in general) γ -law perfect fluid. In section 3, by exploiting the basic relations of the metric approach and the results of paper I we find all the (simply) transitively self-similar SH models i.e. those models which admit a proper Homothetic Vector Field (HVF) without assuming a specific matter content filled the spacetime. From these results we proceed in section 4 and give an analysis of their possible physical interpretation. Due to their sound dynamical significance, we apply the general geometric results only to the case of vacuum and γ -law perfect fluid models. For illustration purposes we reproduce the known vacuum and non-tilted self-similar models of type III, IV, VI_h. In the case of tilted perfect fluids we find a *new exact solution*, representing the *general* self-similar tilted model of type VI_{-1/9}. Finally section 5 contains a summary of all the known self-similar tilted perfect fluid solutions and a brief discussion regarding their importance in the asymptotic behaviour of generic models.

Throughout the following conventions have been used: spatial frame indices are denoted by lower Greek letters $\alpha, \beta, \dots = 1, 2, 3$, lower Latin letters denote spacetime indices $a, b, \dots = 0, 1, 2, 3$ and we use geometrised units such that $8\pi G = c = 1$.

2. Preliminaries

Using the so-called *metric* approach, one can easily derive the explicit form (in local coordinates) of the self-similar metric and the conformally mapped tilted fluid velocity. Due to the fact that the reduced Field Equations (FE) have purely algebraic form, the determination of a specific self-similar vacuum or γ -law perfect fluid model is straightforward. On the other hand the *orthonormal frame* approach is used to write the FE in an autonomous form and the resulting system of decoupled first order differential equations is studied in the *dynamical state space* of the corresponding model [7]. Since every (non-extreme) equilibrium point is represented by an exact self-similar solution the mutual use of the approaches is able to show the existence or not of a self-similar model and to provide with the necessary tools for the description of their future asymptotic dynamics. Therefore and in order the present article to be self-contained as possible we find convenient to give the general setup concerning the essence of both approaches which shall be used frequently in what follows (the reader is referred to [6, 7, 8, 9, 10] for an extensive exposition of the formalism used in this section).

2.1. The metric approach

In Bianchi models the existence of a $\mathcal{G}_3$ Lie algebra of Killing Vector Fields (KVF's) $\mathbf{X}_\alpha$ with 3-dimensional spacelike orbits $\mathcal{C}$ implies the existence of a uniquely defined unit timelike congruence n^a ($n^a n_a = -1$) normal to the spatial foliations $\mathcal{C}$:

$$n_{[a;b]} = 0 = n_{a;b} n^b \Leftrightarrow \frac{1}{2} \mathcal{L}_n g_{ab} \equiv n_{a;b} = \sigma_{ab} + \frac{\theta}{3} h_{ab} \quad (2.1)$$

where $\sigma_{ab}, \theta = n_{a;b}g^{ab}, h_{ab} = g_{ab} + n_a n_b$ are the kinematical quantities associated with the n^a according to the standard kinematical decomposition of an arbitrary timelike congruence [11]. Because n^a is irrotational and geodesic, there exists a time function $t(x^a)$ such that $n^a = \delta_t^a$ i.e. each value of t essentially represents the hypersurfaces $\mathcal{C}$. Therefore local coordinates $\{t, x^\alpha\}$ can be found in which the line element takes the following form:

$$ds^2 = -dt^2 + g_{\alpha\beta}(t)\omega_a^\alpha\omega_b^\beta dx^a dx^b \quad (2.2)$$

where we have employed a basis of invariant vector fields and 1-forms constituting of $\mathbf{e}_\alpha$ and its dual ω^β .

Furthermore if we assume that the spacetime is filled with a perfect fluid having a linear equation of state $\tilde{p} = (\gamma - 1)\tilde{\mu}$, the Ricci tensor is written:

$$R_{ab} = \gamma\tilde{\mu}u_a u_b + \frac{(2 - \gamma)\tilde{\mu}}{2}g_{ab} \quad (2.3)$$

where $\tilde{\mu}, \tilde{p}$ are the energy density and the pressure measured by the observers comoving with the fluid velocity u^a . The tilted (in general) fluid velocity u^a can be decomposed parallel and normal to n^a as follows:

$$\mathbf{u} = \Gamma \mathbf{n} + \Gamma \Delta_\alpha \omega^\alpha = \Gamma \mathbf{n} + \Gamma B^\alpha \mathbf{e}_\alpha \quad (2.4)$$

where $B^\alpha(t), \Delta_\alpha(t)$ are the frame components of the spatial part of the four-velocity u^a and Γ is a smooth function of the time coordinate t satisfying the constraint:

$$\Gamma = (1 - B^\alpha \Delta_\alpha)^{-\frac{1}{2}} \quad (2.5)$$

and $B^\alpha \Delta_\alpha < 1$.

Finally we note that under the assumption of a $\mathcal{H}_4$ Lie algebra of homotheties, it has been shown in paper I that Γ becomes a constant and both the frame components of the metric and the fluid velocity are determined explicitly up to integration constants.

2.2. The orthonormal frame approach

Of particular importance in the study of the asymptotic dynamics of SH models, is the reformulation of the FE as an autonomous system of first order ordinary differential equations. This has been done in [6] where, using the orthonormal frame approach, the FE are written in terms of a set of expansion-normalised variables, by using the dimensionless time parameter:

$$\frac{dt}{d\tau} = \frac{1}{H}, \quad \frac{dH}{d\tau} = -(1 + q)H \quad (2.6)$$

where q, H being the deceleration and expansion (Hubble) parameter respectively of the normal timelike congruence n^a .

The resulting system consists of the evolution equations for the components $\Sigma_{\alpha\beta}$ of the shear tensor of the unit normal vector field n^a , the spatial curvature $A_\alpha, n_{\alpha\beta}$ of the orbits of the G_3 isometry group (due to Jacobi identities) and the spatial components of the tilted fluid velocity v^α .

Using the freedom of a time-dependent spatial rotation, we may choose the orthonormal tetrad to be the eigenframe of $n_{\alpha\beta}$ therefore the contracted form of Jacobi identities $n_{\alpha\beta}A^\beta = 0$ implies:

$$n_{\alpha\beta} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & N_2 & 0 \\ 0 & 0 & N_3 \end{pmatrix}, \quad A_\alpha = A_1 \delta_\alpha^1. \quad (2.7)$$

We note that in types VI_h and VII_h the evolution equations for $n_{\alpha\beta}$ and A^α can be used to express the component A_1 in terms of the curvature variables:

$$A_1^2 = hN_2N_3. \quad (2.8)$$

In addition and following [6] we introduce the shear variables:

$$\Sigma_+ = \frac{1}{2}(\Sigma_{22} + \Sigma_{33}), \quad \Sigma_- = \frac{1}{2\sqrt{3}}(\Sigma_{22} - \Sigma_{33}) \quad (2.9)$$

$$\Sigma_1 = \frac{1}{\sqrt{3}}\Sigma_{23}, \quad \Sigma_3 = \frac{1}{\sqrt{3}}\Sigma_{12}, \quad \tilde{\Sigma}_{13} = \frac{1}{\sqrt{3}}\Sigma_{13} \quad (2.10)$$

where for the sake of simplicity we can drop the tilde.

With these identifications the expansion-normalized variables $\{\Sigma_+, \Sigma_-, \Sigma_1, \Sigma_3, \Sigma_{13}, N_2, N_3, v_\alpha\}$ satisfy the set of evolution equations (A.11)-(A.13) and (A.29) plus the algebraic constraint (A.18) given in [6]. Consequently the state space is a compact subset $\mathcal{D} \subset \mathbf{R}^7$ (the compactness of $\mathcal{D}$ can be shown using the generalised Friedmann equation and the inequality $\Omega > 0$ for the energy density parameter).

We conclude this section by noting that for each Bianchi type, the evolution equations and the algebraic constraints can be determined by specialising them in the standard way:

Type	N_2	N_3	Restrictions
IV	0	N_3	$N_3 > 0$
V	0	0	None
VI _h	N_2	N_3	$N_2 N_3 < 0$
VII _h	N_2	N_3	$N_2 N_3 > 0$
VI _{-1/9}	N_2	N_3	$N_2 N_3 < 0$

3. Solution of the symmetry equations

In this section we will give the general solution of the similarity equations $\mathcal{L}_{\mathbf{H}}g_{ab} = 2\psi g_{ab}$ where $\mathbf{H} = H\mathbf{n} + A^\alpha \mathbf{X}_\alpha$ is the generator of the one-parameter group of homotheties. Using the fact that $[\mathbf{H}, \mathbf{X}_\alpha] = C_{4\alpha}^\beta \mathbf{X}_\beta$ and $[\mathbf{n}, \mathbf{X}_\alpha] = 0$ it follows $H = H(t)$ and $A = A(x^\alpha)$ where $H(t), A(x^\alpha)$ are smooth functions of the spacetime manifold [1]. We note that the exact form of the self-similar SH metrics is *independent* from the physical assumption we made for the matter content of the spacetime. However if we employ a perfect fluid source for the gravitational field, the well known relation $\mathcal{L}_{\mathbf{H}}R_{ab} = \mathcal{L}_{\mathbf{H}}T_{ab} = 0$ implies [12] that the fluid velocity is mapped conformally by the HVF i.e. $\mathcal{L}_{\mathbf{H}}u_a = \psi u_a$.

In what follows the coordinate forms of the KVF's, the invariant basis and its dual are those which appear in [10]. We recall that in types VI_h and VII_h the group parameters h_R and h are related according to $h = -\frac{(h_R+1)^2}{(1-h_R)^2}$ and $h = \frac{h_R^2}{4-h_R^2}$ respectively.

Type III

In contrast with the type VI_{-1/9} models ($h = -1/9 \Leftrightarrow h_R = -2$ or $h_R = -1/2$) in which its "exceptional" dynamical behaviour do not pass to geometry (at least regarding the determination of the corresponding self-similar metrics), type III models ($h = -1 \Leftrightarrow h_R = 0$) cannot be treated simultaneously within the class of type VI_h metrics due to the "singular" property of the value $h_R = 0$ in Jacobi identities.

In local coordinates $\{t, x, y, z\}$ the KVF's $\{\mathbf{X}_\alpha\}$ and the canonical 1-forms $\{\omega^\alpha\}$ are:

$$\mathbf{X}_1 = \partial_y, \quad \mathbf{X}_2 = \partial_z, \quad \mathbf{X}_3 = \partial_x + y\partial_y \quad (3.1)$$

$$\omega^1 = e^{-x}dy, \quad \omega^2 = dz, \quad \omega^3 = dx. \quad (3.2)$$

The non-vanishing structure constants $C_{13}^1 = 1$ and the Jacobi identities of the homothetic Lie algebra $\mathcal{H}_4$ imply that the remaining non-vanishing structure constants $C_{\beta 4}^\alpha$ are either $C_{24}^2 = b$ or $C_{34}^3 = c$. We consider subcases (in order to pertain the generality of the results we avoid to set some group constants equal to 1):

Case A₁ HVF

$$\mathbf{H} = \psi t \partial_t + D_1 \partial_x + b z \partial_z \quad (3.3)$$

Fluid velocity

$$\Delta_1 = v_1 t^{(D_1+\psi)/\psi}, \quad \Delta_2 = v_2 t^{(\psi-b)/\psi}, \quad \Delta_3 = v_3 t \quad (3.4)$$

Metric

$$g_{\alpha\beta} = \begin{pmatrix} c_{11} t^{2(D_1+\psi)/\psi} & c_{12} t^{(D_1+2\psi-b)/\psi} & c_{13} t^{(2\psi+D_1)/\psi} \\ c_{12} t^{(D_1+2\psi-b)/\psi} & c_{22} t^{2(\psi-b)/\psi} & c_{23} t^{(2\psi-b)/\psi} \\ c_{13} t^{(2\psi+D_1)/\psi} & c_{23} t^{(2\psi-b)/\psi} & c_{33} t^2 \end{pmatrix}. \quad (3.5)$$

Case A₂
HVF

$$\mathbf{H} = \psi t \partial_t + D_1 \partial_x + D_2 e^x \partial_y + c x \partial_z \quad (3.6)$$

Fluid velocity

$$\Delta_1 = v_1 t^{(D_1+\psi)/\psi}, \quad \Delta_2 = v_2 t, \quad \Delta_3 = -\frac{D_2 v_1 t^{(D_1+\psi)/\psi}}{D_1} - \frac{c v_2 t \ln t}{\psi} + v_3 t \quad (3.7)$$

Metric

$$g_{\alpha\beta} = \begin{pmatrix} c_{11} t^{2(\psi+D_1)/\psi} & g_{21} & g_{31} \\ c_{12} t^{(2\psi+D_1)/\psi} & c_{22} t^2 & g_{32} \\ t^{(2\psi+D_1)/\psi} \left(c_{13} - \frac{c_{12} c}{\psi} \ln t \right) - \frac{D_2 c_{11} t^{2(\psi+D_1)/\psi}}{D_1} & g_{32} & g_{33} \end{pmatrix} \quad (3.8)$$

where

$$g_{32} = t^2 \left(c_{23} - \frac{c_{22} c \ln t}{\psi} \right) - \frac{c_{12} D_2 t^{(2\psi+D_1)/\psi}}{D_1} \quad (3.9)$$

$$g_{33} = \frac{D_2^2 c_{11}}{D_1^2} t^{2(D_1+\psi)/\psi} + t^{(D_1+2\psi)/\psi} \left(\frac{2D_2 c_{12} c \ln t}{D_1 \psi} - \frac{2D_2 c_{13}}{D_1} \right) + \left[\frac{c^2 c_{22} (\ln t)^2}{\psi^2} - \frac{2c_{23} c \ln t}{\psi} + c_{33} \right] t^2. \quad (3.10)$$

Case A₃**HVF**

$$\mathbf{H} = \psi t \partial_t + D_2 e^x \partial_y + c x \partial_z \quad (3.11)$$

Fluid velocity

$$\Delta_1 = v_1 t, \quad \Delta_2 = v_2 t, \quad \Delta_3 = t \left(v_3 - \frac{D_2 v_1 + c v_2}{\psi} \ln t \right) \quad (3.12)$$

Metric

$$g_{\alpha\beta} = \begin{pmatrix} c_{11} t^2 & g_{21} & g_{31} \\ c_{12} t^2 & c_{22} t^2 & g_{32} \\ t^2 \left(c_{13} - \frac{D_2 c_{11} + c_{12} c}{\psi} \ln t \right) & t^2 \left(c_{23} - \frac{D_2 c_{12} + c_{22} c}{\psi} \ln t \right) & g_{33} \end{pmatrix} \quad (3.13)$$

$$\text{where } g_{33} = t^2 \left[c_{33} + \frac{D_2^2 c_{11} + 2D_2 c_{12} c + c^2 c_{22}}{\psi^2} (\ln t)^2 - \frac{2(D_2 c_{13} + c_{23} c) \ln t}{\psi} \right].$$

Type IV

In type IV models, the KVF's and the dual basis have the following coordinate forms:

$$\mathbf{X}_1 = \partial_y, \quad \mathbf{X}_2 = \partial_z, \quad \mathbf{X}_3 = \partial_x + (y+z)\partial_y + z\partial_z \quad (3.14)$$

$$\omega^1 = e^{-x}dy - xe^{-x}dz, \quad \omega^2 = e^{-x}dz, \quad \omega^3 = dx. \quad (3.15)$$

Since the non-vanishing structure constants are $C_{13}^1 = C_{23}^1 = C_{23}^2 = 1$ a similar analysis shows that the remaining non-vanishing structure constants $C_{\beta 4}^\alpha$ of $\mathcal{H}_4$ are $C_{14}^1 = a$, $C_{24}^2 = a$. Therefore we have the following results:

HVF

$$\mathbf{H} = \psi t \partial_t + D_1 \partial_x + ay \partial_y + az \partial_z \quad (3.16)$$

Fluid velocity

$$\Delta_1 = v_1 t^{(D_1+\psi-a)/\psi}, \quad \Delta_2 = t^{(D_1+\psi-a)/\psi} \left(v_2 + \frac{D_1 v_1 \ln t}{\psi} \right), \quad \Delta_3 = v_3 t \quad (3.17)$$

Metric

$$g_{\alpha\beta} = \begin{pmatrix} c_{11} t^{2(D_1+\psi-a)/\psi} & g_{21} & g_{31} \\ t^{2(D_1+\psi-a)/\psi} \left(c_{12} + \frac{D_1 c_{11} \ln t}{\psi} \right) & g_{22} & g_{32} \\ c_{13} t^{(D_1+2\psi-a)/\psi} & t^{(D_1+2\psi-a)/\psi} \left(c_{23} + \frac{D_1 c_{13} \ln t}{\psi} \right) & c_{33} t^2 \end{pmatrix} \quad (3.18)$$

where

$$g_{22} = t^{2(D_1+\psi-a)/\psi} \left[c_{22} + \frac{2D_1 c_{12} \ln t}{\psi} + \frac{D_1^2 c_{11} (\ln t)^2}{\psi^2} \right]. \quad (3.19)$$

Type V

In this case the KVF's $\{\mathbf{X}_\alpha\}$ and the canonical 1-forms $\{\omega^\alpha\}$ are:

$$\mathbf{X}_1 = \partial_y, \quad \mathbf{X}_2 = \partial_z, \quad \mathbf{X}_3 = \partial_x + y\partial_y + z\partial_z. \quad (3.20)$$

$$\omega^1 = e^{-x}dy, \quad \omega^2 = e^{-x}dz, \quad \omega^3 = dx \quad (3.21)$$

It turns out that the structure of the homothetic algebra is $C_{13}^1 = C_{23}^2 = 1$, $C_{14}^2 = b$, $C_{24}^1 = a$. The analysis of the symmetry equations suggests that we must again consider subcases according to the vanishing of the parameter a :

Case A₁ ($a \neq 0$)**HVF**

$$\mathbf{H} = \psi t \partial_t + D_1 \partial_x + bz \partial_y + ay \partial_z \quad (3.22)$$

Fluid velocity

$$\Delta_1 = v_1 e^{\frac{(D_1 - \sqrt{ab} + \psi) \ln t}{\psi}} + v_2 e^{\frac{(D_1 + \sqrt{ab} + \psi) \ln t}{\psi}}, \quad (3.23)$$

$$\Delta_2 = -\frac{\psi t (\Delta_1)_{,t} - (D_1 + \psi) \Delta_1}{a}, \quad \Delta_3 = v_3 t \quad (3.24)$$

Metric

$$g_{11} = c_{12} e^{\frac{2(D_1 - \sqrt{ab} + \psi) \ln t}{\psi}} + c_{21} e^{\frac{2(D_1 + \sqrt{ab} + \psi) \ln t}{\psi}} + c_{23} t^{\frac{2(D_1 + \psi)}{\psi}} \quad (3.25)$$

$$g_{12} = -\frac{\psi t (g_{11})_{,t} - 2(D_1 + \psi) g_{11}}{2a}, \quad g_{13} = c_{13} e^{\frac{(D_1 - \sqrt{ab} + 2\psi) \ln t}{\psi}} + c_{31} e^{\frac{(D_1 + \sqrt{ab} + 2\psi) \ln t}{\psi}} \quad (3.26)$$

$$g_{22} = \frac{\psi^2 t^2 (g_{11})_{,tt} - \psi t (4D_1 + 3\psi) (g_{11})_{,t} + 2(2D_1^2 + 4D_1\psi - ab + 2\psi^2) g_{11}}{2a^2} \quad (3.27)$$

$$g_{23} = -\frac{\psi t (g_{13})_{,t} - (D_1 + 2\psi) g_{13}}{a}, \quad g_{33} = c_{33} t^2 \quad (3.28)$$

Case A₂**HVF**

$$\mathbf{H} = \psi t \partial_t + D_1 \partial_x + bz \partial_y \quad (3.29)$$

Fluid velocity

$$\Delta_1 = v_1 t^{\frac{D_1+\psi}{\psi}}, \quad \Delta_2 = t^{\frac{D_1+\psi}{\psi}} \left(v_2 - \frac{bv_1 \ln t}{\psi} \right), \quad \Delta_3 = v_3 t \quad (3.30)$$

Metric

$$g_{\alpha\beta} = \begin{pmatrix} c_{11} t^{\frac{2(D_1+\psi)}{\psi}} & t^{\frac{2(D_1+\psi)}{\psi}} \left[c_{22} - \frac{2bc_{12} \ln t}{\psi} + \frac{b^2 c_{11} (\ln t)^2}{\psi^2} \right] & g_{31} \\ t^{\frac{2(D_1+\psi)}{\psi}} \left(c_{12} - \frac{bc_{11} \ln t}{\psi} \right) & t^{\frac{D_1+2\psi}{\psi}} \left(c_{23} - \frac{bc_{13} \ln t}{\psi} \right) & g_{32} \\ c_{13} t^{\frac{D_1+2\psi}{\psi}} & c_{33} t^2 & \end{pmatrix}. \quad (3.31)$$

Type VI_{h_R}

The KVF's $\{\mathbf{X}_\alpha\}$ and the canonical 1-forms $\{\omega^\alpha\}$ are:

$$\mathbf{X}_1 = \partial_y, \quad \mathbf{X}_2 = \partial_z, \quad \mathbf{X}_3 = \partial_x + y \partial_y + h_R z \partial_z. \quad (3.32)$$

$$\omega^1 = e^{-x} dy, \quad \omega^2 = e^{-h_R x} dz, \quad \omega^3 = dx \quad (3.33)$$

where $h_R \neq 0$ and $C_{13}^1 = h_R^{-1} C_{23}^2 = 1$. Hence the remaining non-vanishing structure constants $C_{\beta 4}^\alpha$ are $C_{14}^1 = a$, $C_{24}^2 = b$. Using these results we get:

HVF

$$\mathbf{H} = \psi t \partial_t + D_1 \partial_x + ay \partial_y + bz \partial_z \quad (3.34)$$

Fluid velocity

$$\Delta_1 = v_1 t^{(D_1+\psi-a)/\psi}, \quad \Delta_2 = v_2 t^{(D_1 h_R + \psi - b)/\psi}, \quad \Delta_3 = v_3 t \quad (3.35)$$

Metric

$$g_{\alpha\beta} = \begin{pmatrix} c_{11} t^{2(D_1+\psi-a)/\psi} & c_{22} t^{2(D_1 h_R + \psi - b)/\psi} & g_{31} \\ c_{12} t^{[D_1(h_R+1)+2\psi-a-b]/\psi} & c_{23} t^{(D_1 h_R + 2\psi - b)/\psi} & g_{32} \\ c_{13} t^{(D_1+2\psi-a)/\psi} & c_{33} t^2 & \end{pmatrix}. \quad (3.36)$$

Type VII_{h_R}

Finally in type VII_{h_R} models we have:

$$\mathbf{X}_1 = \partial_y, \quad \mathbf{X}_2 = \partial_z, \quad \mathbf{X}_3 = \partial_x - z \partial_y + (y + h_R z) \partial_z. \quad (3.37)$$

$$\omega^1 = \left(A_1 - \frac{h_R}{2} A_2 \right) dy - A_2 dz, \quad \omega^2 = A_2 dy + \left(A_1 + \frac{h_R}{2} A_2 \right) dz, \quad \omega^3 = dx \quad (3.38)$$

where:

$$A_1 = e^{-\frac{h_R}{2}x} \cos wx, \quad A_2 = -\frac{1}{w} e^{-\frac{h_R}{2}x} \sin wx, \quad w = \sqrt{\frac{4 - h_R^2}{4}}. \quad (3.39)$$

The non-vanishing structure constants are ($h_R^2 < 4$) $C_{13}^2 = -C_{23}^1 = h_R^{-1} C_{23}^2 = 1$ leading to the following expressions:

$$C_{14}^1 = a, \quad C_{24}^2 = a. \quad (3.40)$$

HVF

$$\mathbf{H} = \psi t \partial_t + D_1 \partial_x + ay \partial_y + az \partial_z \quad (3.41)$$

Fluid velocity

$$\Delta_1 = t^{p_1/2\psi} \left[v_{12} \cos \left(\frac{D_1 w}{\psi} \ln t \right) + v_{21} \sin \left(\frac{D_1 w}{\psi} \ln t \right) \right], \quad (3.42)$$

$$\Delta_2 = \frac{\psi t (\Delta_1)_{,t} + (a - \psi) \Delta_1}{D_1}, \quad \Delta_3 = v_3 t. \quad (3.43)$$

Metric

$$g_{11} = - \frac{t^{p_1/\psi} \left\{ 16 D_1 E_1 w^3 p_1^2 \cos \left(\frac{2 D_1 w \ln t}{\psi} \right) + 4 E_2 w^2 p_1 \sin \left(\frac{2 D_1 w \ln t}{\psi} \right) \right\}}{64 D_1^3 w^5 p_1^2} \quad (3.44)$$

$$g_{12} = \frac{\psi t (g_{11})_{,t} + 2(a - \psi) g_{11}}{2 D_1} \quad (3.45)$$

$$g_{13} = t^{(p_1+2\psi)/2\psi} \left[c_{13} \cos \left(\frac{D_1 w \ln t}{\psi} \right) + c_{31} \sin \left(\frac{D_1 w \ln t}{\psi} \right) \right] \quad (3.46)$$

$$g_{22} = \frac{\psi^2 t^2 (g_{11})_{,tt} - \psi t (g_{11})_{,t} (p_1 + \psi - 2a) + 2 [D_1^2 + (\psi - a) p_1] g_{11}}{2 D_1^2} \quad (3.47)$$

$$g_{23} = \frac{\psi t (g_{13})_{,t} + (a - 2\psi) g_{13}}{D_1}, \quad g_{33} = c_{33} t^2 \quad (3.48)$$

where E_1, E_2 are constants and we have set $p_1 = D_1 h_R + 2(\psi - a)$.

4. Exact Solutions

The well known feature of self-similarity to reduce the FE to a purely algebraic form, makes the determination of vacuum and perfect fluid models straightforward. Following the methodology of paper I and for illustration purposes, we reproduce all the known vacuum and non-tilted perfect fluid solutions. In addition we find the *general* tilted perfect fluid model for the exceptional class VI_{-1/9}. For completeness we also give, for each Bianchi type, the extreme[‡] tilted equilibrium points of their state space.

Vacuum plane wave models of type III

This exact vacuum solution can be found from the case A₁ of section 2 and has been given in [13]. The HVF is $\mathbf{H} = \psi t \partial_t + D_1 \partial_x + bz \partial_z$ and the constants $c_{\alpha\beta}, \psi, b$ are:

$$c_{13} = c_{23} = 0, \quad c_{11} = \frac{c_{12}^2 (D_1 - b)}{2 c_{22} D_1}, \quad c_{33} = \frac{b^2}{(D_1 + b)^2}, \quad \psi = b \quad (4.1)$$

[‡] We recall that an extreme tilted equilibrium point satisfies $v^\alpha v_\alpha = 1$.

where without loss of generality we can set $b = 1$.

We note that the above spacetime is algebraically special (Petrov type N) since it admits the *gradient null KVF* $l^a = e^{-bx/(D_1+b)} (\partial_t + \frac{D_1+b}{bt} \partial_x)$ which essentially represents the repeated principal null direction of the Weyl tensor.

Non-tilted perfect fluid models of type III

It is a special case of the non-tilted perfect fluid solution found by Collins [14] for $h = -1$. The HVF is $\mathbf{H} = \psi t \partial_t + bz \partial_z$ and the constants $c_{\alpha\beta}, \psi, b$ are given by:

$$\begin{aligned} c_{13} = c_{23} = c_{12} = 0, \quad c_{11} = c_{22} = 1 \\ c_{33} = \frac{\gamma^2}{(2-\gamma)(3\gamma-2)}, \quad \mu = \frac{4(1-\gamma)}{\gamma^2 t^2}, \quad \psi = \frac{b\gamma}{3\gamma-2}. \end{aligned} \quad (4.2)$$

Tilted perfect fluid models of type III

Using the results of section 2 we can show that there is no physically acceptable tilted perfect solution of type III. We note that this conclusion can be also confirmed from the set of the tilted equilibrium points of the dynamical state space given in [15].

Vacuum plane wave models of type IV

From the self-similar metrics (3.18) we can find the general type IV vacuum solution, first given in [16]. In this case the HVF is $\mathbf{H} = \psi t \partial_t + D_1 \partial_x + ay \partial_y + az \partial_z$ or, by performing a suitable change of the basis of the homothetic Lie algebra, $\mathbf{H} = \psi t \partial_t + (D_1 - a) \partial_x - az \partial_y$. In addition the constants $c_{\alpha\beta}, \psi$ are given by:

$$c_{13} = c_{23} = 0, \quad c_{22} = \frac{D_1(c_{11}^2 - 4c_{12}^2) + 4ac_{12}^2}{4c_{11}(a - D_1)}, \quad c_{33} = \frac{a^2}{D_1^2}, \quad \psi = a. \quad (4.3)$$

It can be verified that the above one parameter family of models is also algebraically special and admits the *gradient null KVF* $l^a = e^{-ax/D_1} (\partial_t + \frac{D_1}{at} \partial_x)$.

Non-tilted and tilted perfect fluid models of type IV

It is found that no orthogonal or tilted perfect fluid self-similar models of type IV exist. Nevertheless there exists the following extreme tilted equilibrium point [17]:

$$\begin{aligned} N_2 = 0, \quad N_3 = N_3, \quad v^\alpha v_\alpha = 1, \\ \Sigma^2 = \frac{N_3^2 + 3q^2}{12}, \quad \Sigma_3 = \Sigma_{13} = \Sigma_- = 0, \quad \Sigma_+ = -\frac{q}{2}, \\ \Sigma_1 = \frac{\sqrt{3}N_3}{6}, \quad v_1 = 1, \quad A_1 = \frac{2-q}{2} \\ \Omega = \frac{3q(2-q) - N_3^2}{6}, \quad 0 < \gamma < 2. \end{aligned}$$

Models of type V

The analysis shown that no self-similar vacuum or perfect fluid models exist for type V. This result can also be proved by using the set of evolution equations given in [6] (see also [15]). However there exists the following extreme tilted equilibrium point [18, 19]:

$$N_2 = N_3 = 0, \quad v^\alpha v_\alpha = 1,$$

$$\Sigma^2 = (A_1 - 1)^2, \quad \Sigma_3 = \Sigma_{13} = \Sigma_1 = \Sigma_- = 0, \quad \Sigma_+ = -\frac{q}{2},$$

$$v_1 = 1, \quad A_1 = \frac{2-q}{2}, \quad \Omega = 2A_1(1 - A_1), \quad 0 < \gamma < 2.$$

Vacuum plane wave models of type VI_{h_R}

In this case the HVF is $\mathbf{H} = \psi t \partial_t + D_1 \partial_x + ay \partial_y + bz \partial_z$ and the constants $c_{\alpha\beta}, \psi$ are given by [13]:

$$c_{13} = c_{23} = 0, \quad c_{11} = \frac{c_{12}^2 (h_R + 1) [D_1 (h_R + 1) - a - b]}{2c_{22} [D_1 (h_R^2 + 1) - a - bh_R]} \quad (4.4)$$

$$c_{33} = \frac{(ah_R - b)^2}{[D_1 (h_R - 1) + a - b]^2}, \quad \psi = \frac{ah_R - b}{h_R - 1}.$$

The *gradient null KVF* is $l^a = e^{-\frac{ah_R - b}{D_1(h_R - 1) + a - b}x} \left[\partial_t + \frac{D_1(h_R - 1) + a - b}{(ah_R - b)t} \partial_x \right]$.

As we have mentioned in section 2, in order to determine the corresponding self-similar solution for the exceptional type VI_{-1/9} models we can use the geometric results (3.34)-(3.36) and solve the FE by setting $h_R = -2$ or $h_R = -1/2$ (for these values of the group parameter the 01 or 02 component of (2.3) vanish and essentially represents the "exceptional" behaviour of the VI_{-1/9} models in terms of the FE). The resulting solution has been first given by Robinson and Trautman (RT) [20] (see also [13]) and has the following form ($c_{13} \neq 0$):

$$c_{12} = c_{23} = 0, \quad c_{33} = \frac{75}{8}, \quad c_{11} = \frac{24c_{13}^2}{125}, \quad D_1 = -\frac{b}{3}, \quad \psi = \frac{5b}{6}, \quad a = \frac{b}{3}. \quad (4.5)$$

Non-tilted perfect fluid models of type VI_{h_R}

The family of non-tilted perfect fluid models has been found by Collins [14]. The HVF is given by equation (3.34) and the constants $c_{\alpha\beta}, \psi, a, b$ are:

$$c_{13} = c_{23} = c_{12} = 0, \quad c_{11} = c_{22} = 1, \quad c_{33} = \frac{\gamma^2 (h_R - 1)^2}{(2 - \gamma)(3\gamma - 2)}$$

$$\mu = -\frac{4[h_R^2(\gamma - 1) + h_R\gamma + \gamma - 1]}{\gamma^2 (h_R - 1)^2 t^2}, \quad \psi = \frac{\gamma(D_1 - a)(h_R - 1)}{h_R(2 - 3\gamma)} \quad (4.6)$$

$$b = \frac{D_1(h_R^2 + 1) - a}{h_R}.$$

The exceptional type VI_{-1/9} non-tilted perfect fluid solution can be found similarly by setting $h_R = -2$ (or $h_R = -1/2$) and has been given in [21]:

$$c_{12} = c_{23} = 0, \quad c_{33} = \frac{75}{8}, \quad \psi = \frac{5b}{2}, \quad D_1 = 0, \quad a = 2b \quad (4.7)$$

$$\mu = \frac{9(125c_{11} - 24c_{13}^2)}{25(75c_{11} - 8c_{13}^2)t^2}, \quad \gamma = \frac{10}{9}.$$

Tilted perfect fluid models of type VI_{h_R}

It is convenient to employ the constants p_1, p_2 :

$$D_1 = a + p_1\psi - 2\psi, \quad a = bh_R - \psi(p_1 + p_2 - 4). \quad (4.8)$$

The frame components of the metric and the fluid velocity become:

$$g_{\alpha\beta} = \begin{pmatrix} c_{11}t^{2(p_1-1)} & g_{21} & g_{31} \\ c_{12}t^{[b(h_R^2-1)+h_R\psi(2-p_2)+p_1\psi]/\psi} & c_{22}t^{2[b(h_R^2-1)+h_R\psi(2-p_2)+\psi]/\psi} & g_{32} \\ c_{13}t^{p_1} & c_{23}t^{[b(h_R^2-1)+h_R\psi(2-p_2)+2\psi]/\psi} & c_{33}t^2 \end{pmatrix} \quad (4.9)$$

$$\Delta_1 = v_1 t^{p_1-1}, \quad \Delta_2 = v_2 t^{[b(h_R^2-1)+h_R\psi(2-p_2)+\psi]/\psi}, \quad \Delta_3 = v_3 t \quad (4.10)$$

whereas the HVF assumes the form:

$$\mathbf{H} = \psi t \partial_t + [bh_R - \psi(p_2 - 2)] \partial_x + [bh_R - \psi(p_1 + p_2 - 4)] y \partial_y + bz \partial_z. \quad (4.11)$$

For the case $h_R \neq -2$ (or $h_R \neq -1/2$) the general self-similar solution has been found recently [15]. Here we give the corresponding *general solution* for the type VI_{-1/9} models.

We define the parameter s according to:

$$\gamma = \frac{2}{2s+1}. \quad (4.12)$$

Then, the 00-conservation equation implies that the constant p_1 is given by:

$$p_1 = \frac{2(6s+1) - p_2}{4}. \quad (4.13)$$

The various integration constants are:

$$c_{12} = c_{13} = v_1 = 0, \quad \psi = -\frac{2b}{p_2 - 2} \quad (4.14)$$

$$c_{22} = \frac{c_{23}^2(p_2-2)^2[2(3s-1)-p_2] \{-9p_2^2+4p_2(3-2s)+624s^2-208s+12\}}{96s[25p_2^2-20p_2(6s+1)+4(6s+1)^2]}$$

$$c_{33} = \frac{48[3p_2^2+4p_2(8s-3)-4(12s^2+8s-3)]}{(p_2-2)^2[9p_2^2+4p_2(2s-3)-4(156s^2-52s+3)]}$$

$$\Gamma^2 = -\frac{s(5p_2-12s-2)[2(3s-1)-p_2] \{-9p_2^2+4p_2(3-2s)+624s^2-208s+12\}}{2[p_2+2(2s-1)][p_2^2(31s-6)+4p_2s(78s-17)-4(1476s^3-660s^2+101s-6)]} \quad (4.15)$$

$$v_3 = -\frac{36(p_2+4s-2)[p_2+2(2s-1)]}{9p_2^3+2p_2^2(4s-15)-12p_2(52s^2-16s-1)+8(156s^2-52s+3)}$$

$$v_2 = \frac{3c_{23}[p_2^2+4p_2(s-1)-4(2s-1)]}{4[2(6s+1)-5p_2]}.$$

The constant p_2 is related with the "state parameter" s according to:

$$p_2 = \frac{2[-6\sqrt{36s^4-204s^3+133s^2-22s+1}|4s-1|+s(336s^2-214s+19)]}{160s^2-91s+6} \quad (4.16)$$

and the energy density of the model is:

$$\tilde{\mu} = \frac{(2s+1)[p_2^2(31s-6)+4p_2s(78s-17)-4(1476s^3-660s^2+101s-6)]}{144s(p_2-12s+2)t^2} \quad (4.17)$$

In order to ensure the positivity of the energy density and the real values of the constant p_2 (together with the overall signature of the metric) this new exact solution is only defined when:

$$s \in \left(\frac{1}{6}, \frac{1}{4}\right) \Leftrightarrow \gamma \in \left(\frac{4}{3}, \frac{3}{2}\right). \quad (4.18)$$

It is interesting to note that, similarly with the types VI_0 and VI_h , the above *rotating* cosmological model admits the *hypersurface orthogonal* KVF $\mathbf{X}_2 = \partial_z$ and belongs to the subclass $n_\alpha^\alpha = 0$. In fact the present solution arises as an equilibrium point of the tilted perfect fluid Bianchi type $\text{VI}_{-1/9}$ dynamical state space which we denote it as $\mathcal{E}$ and has the following kinematical and dynamical quantities:

$$\begin{aligned}
N_3 &= -\sqrt{-\frac{3[q(3\gamma-6)+7\gamma-6][q^2(4\gamma^2-4\gamma-12)-2q(3\gamma^2-22\gamma+12)-37\gamma^2+48\gamma-12]}{16\gamma^2[q(\gamma-2)+2(\gamma-1)]}} \\
N_2 &= -N_3, \quad \Sigma_- = 0, \quad \Sigma_+ = -\frac{q}{2}, \quad v_2 = -v_3 \\
\Sigma_1 &= -\frac{\sqrt{3}[q(5\gamma-12)+2(7\gamma-6)]}{6\gamma}, \quad \Sigma_3 = -\frac{\sqrt{6(2-\gamma)}[2q^2(\gamma-3)+3q(3\gamma-4)+7\gamma-6]}{12\gamma\sqrt{(q+1)[q(\gamma-2)+2(\gamma-1)]}} \\
\Sigma_{13} &= \frac{\sqrt{6(2-\gamma)}[2q^3(\gamma-3)+q^2(11\gamma-18)+2q(8\gamma-9)+7\gamma-6]}{12\gamma\sqrt{(q+1)^3[q(\gamma-2)+2(\gamma-1)]}} \\
v_1 &= \frac{3(2q-3\gamma+2)}{2N_3\gamma}, \quad v_3 = -\frac{3\sqrt{2(2-\gamma)(q+1)}(2q-3\gamma+2)}{8N_3\gamma\sqrt{q(\gamma-2)+2(\gamma-1)}}, \quad v^2 = \frac{9(2q-3\gamma+2)^2[3q(\gamma-2)+7\gamma-6]}{16N_3^2\gamma^2[q(\gamma-2)+2(\gamma-1)]} \\
\Omega &= \frac{[q(3\gamma-6)+7\gamma-6][2q(\gamma-3)+7\gamma-6][4q^2(4-\gamma)+2q(8-15\gamma)+27\gamma^2-26\gamma]}{12\gamma^2(2q-3\gamma+2)[q(\gamma-2)+2(\gamma-1)]} \\
\Sigma^2 &= \frac{[3q(\gamma-2)+7\gamma-6][4q^2(2\gamma^2-8\gamma+9)+4q(9\gamma^2-29\gamma+18)+49\gamma^2-84\gamma+36]}{12\gamma^2[q(\gamma-2)+2(\gamma-1)]} \\
q &= \frac{\gamma\sqrt{(\gamma-1)(73\gamma^3-253\gamma^2+240\gamma-36)}|3\gamma-4|-2(\gamma-1)(33\gamma^3-179\gamma^2+294\gamma-144)}{6(\gamma-1)(\gamma-2)(3\gamma^2-19\gamma+24)}
\end{aligned}$$

We also note that there exists the following extreme tilted equilibrium point [19]:

$$\begin{aligned}
N_2 &= N_2, \quad N_3 = N_3, \quad v^\alpha v_\alpha = 1, \\
\Sigma^2 &= \frac{3(N_3 - N_2)^2 - 4(N_2 N_3 + 6\sqrt{-N_2 N_3} - 9)}{36}, \quad \Sigma_3 = \Sigma_{13} = \Sigma_- = 0, \\
\Sigma_1 &= \frac{\sqrt{3}(N_3 - N_2)}{6}, \quad \Sigma_+ = -\frac{q}{2}, \quad q = \frac{2(3 - \sqrt{-N_2 N_3})}{3}, \quad v_1 = 1, \\
\Omega &= -\frac{3(N_3 - N_2)^2 - 4(N_2 N_3 + 3\sqrt{-N_2 N_3})}{18}, \quad 0 < \gamma < 2.
\end{aligned}$$

Models of type VII_{h_R}

The question of the existence of tilted perfect type VII_h models is notoriously difficult to answer due to the complexity of the self-similar metrics (3.44)-(3.48). In fact the resulting FE are difficult to handle analytically even to the case of expressing the known VII_h vacuum solution (see e.g. [7] page 192 for an elegant form of this solution and [22] for a complete study of its stability properties against vorticity, shear and Weyl curvature perturbations). Nevertheless, using the set of evolution equations we can show that the assumption of the existence of a proper HVF is incompatible with tilted perfect fluid models and that type VII_h models possess the following extreme tilted equilibrium point (this equilibrium point has been also given in [17]):

$$\begin{aligned}
N_2 &= N_2, \quad N_3 = N_3, \quad v^\alpha v_\alpha = 1, \\
\Sigma^2 &= \frac{(N_3 - N_2)^2 + 12(hN_2 N_3 - 2\sqrt{hN_2 N_3} + 1)}{12}, \quad \Sigma_3 = \Sigma_{13} = \Sigma_- = 0,
\end{aligned}$$

$$\Sigma_1 = \frac{\sqrt{3}(N_3 - N_2)}{6}, \quad \Sigma_+ = -\frac{q}{2}, \quad q = 2(1 - \sqrt{hN_2N_3}), \quad v_1 = 1,$$

$$\Omega = -\frac{(N_3 - N_2)^2 + 12(hN_2N_3 - \sqrt{hN_2N_3})}{6}, \quad 0 < \gamma < 2.$$

5. Discussion

The main concern of papers I and II can be regarded having two main branches:

- a) a geometric nature concerning the determination of all the SH geometries restricted *only* by the requirement of admitting a proper HVF and,
- b) the physical implications of the general geometric results by finding, whenever they exist, the corresponding vacuum or perfect fluid models.

The underlying importance of a) is that we have not incorporated a specific form for the matter fields filled the spacetime therefore we can associate the self-similar metrics with more general matter configurations with a view to analyse their physical significance. On the other hand the well established aspect of the SH equilibrium points as past or future attractors for general vacuum or perfect fluid models necessitates the knowledge of all the self-similar equilibrium points in order to gain deeper insight into their asymptotic dynamics. This was accomplished in the series of articles I, II, [3, 4, 15] in which the *complete set* of tilted perfect fluid self-similar models have been found. For convenience we collect them in table 1 together with the corresponding references.

As an immediate application of the geometric results of section 3, we reproduced the known exact vacuum and non-tilted solutions for the Bianchi types of class B and we have found the *general* form of the self-similar tilted type $VI_{-1/9}$ models. As we have seen in section 4 this *new* exact solution arise as equilibrium point in the $VI_{-1/9}$ state space and shares many of the common properties of type VI_0 and VI_h ($h \neq -1/9$) models and important differences as well. For example the stability analysis of vacuum and non-tilted equilibrium points [25] has shown that the Collins $VI_{-1/9}$ solution is stable for $\gamma \in (\frac{2}{3}, \frac{10}{9})$. At the arc of equilibrium points $\gamma = \frac{10}{9}$ [24] there is an exchange of stability with the RT vacuum solution [8, 20] which is stable whenever $\gamma \in (\frac{10}{9}, \frac{4}{3})$. As a consequence we expected that tilted models will be future dominated against the RT model when $\gamma > \frac{4}{3}$. Indeed the equilibrium point $\mathcal{E}$ is found to have a five dimensional stable manifold for $\gamma \in (\frac{4}{3}, \frac{3}{2})$. However it can be shown that, at least for the case where $n_\alpha^\alpha = 0$, there exist always two unstable modes in the restricted state space. One can made a step further and conjecture that for the type VI_h and $VI_{-1/9}$ models the values $\gamma = \frac{2(3+\sqrt{-h})}{5+3\sqrt{h}}$ and $\gamma = \frac{4}{3}$ respectively represent line bifurcations similarly to the type VI_0 models with $\gamma = \frac{6}{5}$. However no such bifurcations are found in both models.

Furthermore, for the remaining types, the mutual use of the metric and orthonormal frame approaches guarantees that tilted perfect fluid models of types III, IV, V and VII_h cannot admit a proper HVF indicating that these models either may not be asymptotically self-similar (a preliminary analysis suggests that this behaviour occurs in type V models) or may be extreme tilted at late times (see [17] for a thorough discussion on the asymptotic dynamics of type IV models).

We conclude by noting that an interesting aspect of the analysis presented in papers I and II is that vacuum and tilted perfect fluid SH self-similar models exhibit further geometric constraints coming from the existence of covariantly constant null vector fields or a hypersurface orthogonal KVF. Therefore we believe that it will be interesting to study how the assumptions of the self-similarity and a specific dynamic description lead to extra restrictions on the geometry of the SH models.

Acknowledgments

The author wishes to acknowledge the financial support of Ministry of National Education and Religious Affairs through the research program "Pythagoras", Grant No 70-03-7310.

Table 1. This table contains all the self-similar tilted perfect fluid SH models and the relevant range of the state parameter γ .

Type	Self-similar Model	Reference
I	$\nexists$	[23]
II	$\gamma \in (\frac{10}{7}, 2)$	[2]
VI ₀	$\gamma \in [\frac{6}{5}, \frac{3}{2})$	[4, 5]
VII ₀	$\nexists$	[1]
VIII	$\nexists$	[3]
IX	$\nexists$	[3]
III	$\nexists$	[15]
IV	$\nexists$	paper II
V	$\nexists$	paper II
VI _h	$-1 < h < -\frac{1}{9} \Rightarrow 1 < \gamma < \frac{2(3+\sqrt{-h})}{5+3\sqrt{-h}}$	[15]
	$-\frac{1}{9} < h < 0 \Rightarrow \frac{2(3+\sqrt{-h})}{5+3\sqrt{-h}} < \gamma < \frac{3}{2}$	
VII _h	$\nexists$	paper II
VI _{-1/9}	$\gamma \in (\frac{4}{3}, \frac{3}{2})$	paper II

- [1] Apostolopoulos P S, *Self-similar Bianchi models: I. Class A models*, 2003 Class. Quantum Grav. **20** 3371-3384.
- [2] Hewitt C G, *An exact tilted Bianchi II cosmology*, 1991 Class. Quantum Grav. **8** L109-L114.
- [3] Apostolopoulos P S and Tsamparlis M, *Self-similar Bianchi type VIII and IX models*, 2003 Gen. Relativ. Grav. **35** 2051-2056.
- [4] Apostolopoulos P S, *On tilted perfect fluid Bianchi type VI₀ self-similar models*, 2004 Gen. Relativ. Grav. **36** 1939-1945 (see also **preprint gr-qc/0310033**).
- [5] Rosquist K and Jantzen R T, *Exact power law solutions of the Einstein equations*, 1985 Phys. Lett. **107A** 29-32.
- [6] Hewitt C G, Bridson R and Wainwright J, *The asymptotic regimes of tilted Bianchi II cosmologies*, 2001 Gen. Relativ. Grav. **33** 65-94.
- [7] Wainwright J and Ellis G F R (Editors), *Dynamical Systems in Cosmology* (Cambridge University Press, Cambridge 1997).
- [8] Ellis G F R and MacCallum M A H, *A class of homogeneous cosmological models*, 1969 Commun. Math. Phys. **12** 108-141.
- [9] MacCallum M A H, *Anisotropic and inhomogeneous relativistic cosmologies*, General Relativity, Editors. S. W. Hawking and W. Israel (Cambridge University Press, Cambridge 1979).
- [10] Ryan M P Jr and Shepley L C, *Homogeneous Relativistic Cosmologies*, (Princeton University Press, 1975).
- [11] Ellis G F R, *Cargèse Lectures in Physics, Vol. 6*, Ed. E Schatzman (Gordon and Breach, New York 1973).
- [12] Carot J and Sintès A M, *Homothetic perfect fluid spacetimes*, 1997 Class. Quantum Grav. **14** 1183-1205.
- [13] Hsu L and Wainwright J, *Self-similar spatially homogeneous cosmologies: orthogonal perfect fluid and vacuum solutions*, 1986 Class. Quantum Grav. **3** 1105-1124.
- [14] Collins C B, *More qualitative cosmology*, 1971 Commun. Math. Phys. **23** 137-156.
- [15] Apostolopoulos P S, *Equilibrium points of the tilted perfect fluid Bianchi type VI_h state space*, 2004 to appear in Gen. Relativ. Grav. (**preprint gr-qc/0407040**).
- [16] Harvey A and Tsoubelis D, *Exact Bianchi IV cosmological model*, 1977 Phys. Rev. D **15** 2734-2737.
- [17] Hervik S, van den Hoogen R, Coley A, *Future asymptotic behaviour of tilted Bianchi models of type IV and VII_h*, 2005 Class. Quantum Grav. **22** 607-633 (**preprint gr-qc/0409106**).
- [18] Hewitt C G and Wainwright J, *Dynamical systems approach to tilted Bianchi cosmologies: Irrotational models of type V*, 1992 Phys. Rev. D **46** 4242-4252.
- [19] Coley A A and Hervik S, *A dynamical systems approach to the tilted Bianchi models of solvable type*, 2005 Class. Quantum Grav. **22** 579-605 (**preprint gr-qc/0409100**).
- [20] Robinson I and Trautman A, *Some spherical gravitational waves in general relativity*, 1962 Proc. Royal Soc. London A **265** 463-473.
- [21] Wainwright J, *Power law singularities in orthogonally spatially homogeneous cosmologies*, 1984 Gen. Relativ. Grav.

- 16** 657-674.
- [22] Barrow J and Tsagas C, *Structure and stability of the Lukash plane-wave spacetime*, 2004 (**preprint gr-qc/0411070**).
 - [23] King A R and Ellis G F R, *Tilted Homogeneous Cosmological Models*, 1973 Commun. Math. Phys. **31** 209-242.
 - [24] Hewitt C G, Horwood J T and Wainwright J, *Asymptotic dynamics of the exceptional Bianchi cosmologies*, 2003 Class. Quantum Grav. **20** 1743-1756.
 - [25] Barrow J D and Hervik S, *The future of tilted Bianchi universes*, 2003 Class. Quantum Grav. **20** 2841-2854.